

National Core Indicators®--Intellectual and Developmental Disabilities 2020-21 In-Person Survey (IPS) Report

Medication

Medications are used effectively and appropriately.

What to know about the presentation of Medication data:



Charts with un-collapsed weighted NCI average

Here's what [collapsed](#) data are



State and weighted NCI averages

The number of respondents (N) is not weighted

Here's what [weighting](#) data means for comparisons



DATA CAUTION!

Many states' data have a large amount of missing data or data recorded as “don’t know”

Read about what this means [here](#)

Medication

Table 1. Takes at least one medication for mood, anxiety, psychotic disorder and/or behavioral challenge (information may have been obtained through state records)	7
Table 2. Takes medication for mood, anxiety, and/or psychotic disorders (information may have been obtained through state records)	9
Table 2B. Takes medication for mood, anxiety, and/or psychotic disorders (<i>includes missing and “don’t know” responses</i> ; information may have been obtained through state records)	10
Table 3. Number of medications taken for at least one of the following: mood disorders, anxiety, or psychotic disorders if taking at least one medication for this purpose (information may have been obtained through state records)	12
Table 3B. Number of medications taken for at least one of the following: mood disorders, anxiety, or psychotic disorders if taking at least one medication for this purpose (<i>includes missing and “don’t know” responses</i> ; information may have been obtained through state records)	13
Table 4. Takes medication for behavior challenges (information may have been obtained through state records) .	15
Table 4B. Takes medication for behavior challenges (<i>includes missing and “don’t know” responses</i> ; information may have been obtained through state records)	16
Table 5. Number of medications taken for behavior challenges if taking at least one for this purpose (information may have been obtained through state records)	18
Table 5B. Number of medications taken for behavior challenges if taking at least one for this purpose (<i>includes missing and “don’t know” responses</i> ; information may have been obtained through state records).....	19
Table 6. Has a behavior plan (information may have been obtained through state records)	21
Table 6B. Has a behavior plan (<i>includes missing and “don’t know” responses</i> ; information may have been obtained through state records)	22
Table 7. Has a behavior plan (among those who take medication for a behavior challenge; information may have been obtained through state records).....	24
Table 6B. Has a behavior plan among those who take medication for a behavior challenge (<i>includes missing and “don’t know” responses</i> ; information may have been obtained through state records)	25

Where do these data come from?

Some of the information in this section comes directly from the person receiving services, or for some select questions, from a “proxy” respondent. Others are drawn from administrative sources; these instances are noted in the table titles.

Read more about why we allow proxy respondents for select questions in the IPS in the Overview chapter of this report.

Items in this chapter are a part of the NCI Health, Welfare, and Rights domain (Medication topic area) and comprise the “Medication” sub-domain (a specific area of concern within the domain). The Medication subdomain includes items that help determine whether “medications are managed effectively and appropriately”.

Read more about NCI domains and subdomains in the Overview chapter or this report.

What are “collapsed” data?

Many questions in the IPS allow respondents to answer a question with “middle” response like “maybe”, “sometimes” or “in-between”. The tables in this section present collapsed data. That means that two or more response options are grouped together if they are considered to reflect a positive outcome. The table titles explain which options were grouped.

To see the breakout of responses for each option, please see Appendix C.

What is weighting and why do we do it?

The NCI-IDD averages are “weighted” to reflect the states’ relative population and sample sizes. We created the weights using the state’s number of valid surveys and its total survey-eligible population. This way, a state that provides services to a larger number of people but uses a sample similar in size to other states has a greater influence on the overall NCI-IDD average (that is, its contribution is *proportional to its service population*).

In this survey year, CA intentionally oversampled by geographical regions to get more focused insights, so their weights are calculated slightly differently (using the number of completed surveys and service

population sizes in each subpopulation or region). You can read more about this in the Methodology section of the Overview chapter.

The number of respondents (N) for each question *is not* weighted.

DATA CAUTION!

Several states had large amounts of missing data or data recorded as “don’t know.” Where 25% or more of a state’s sample had “don’t know” or missing responses, we used an asterisk (*) to indicate that state’s data should be interpreted with caution as it may not accurately represent the sample or service population.

Note on WI data – for all respondents, WI was unable to collect several data points from administrative records for questions in the Background Information section. Additional data points from the Background Information section were unable to be collected for those respondents who self-direct their services related to certain demographics, health, medication, and wellness.

Presentation of state and weighted NCI-IDD averages

Averages

For the most part, the state outcomes are shown in descending order of percentage and are categorized as statistically significantly above the NCI-IDD Average, within the range of the NCI-IDD Average, or significantly below the NCI-IDD Average. Statistical significance is taken to be at or below the .01 level. For those states that fall within the NCI-IDD Average range, their ‘always’ or ‘yes’ response was not statistically different from the NCI-IDD Average. Other tables show the states listed in alphabetical order and are not tested for significance.

Tables that show significance also break out average results for each state by the type of residence: ICF/IID or other institutional settings, group residence, own home, parent/relative’s home, foster or host home. Figure A on the following page shows, by state, the percentage of people surveyed in each residence type. Please note though that:

- Significance testing was not performed on the breakouts by residence

- The number of people in each residential setting is often too small to allow for valid state-to-state comparisons
- And in many cases, statistically valid conclusions cannot be drawn about differences in results between residence types

Still, we provide this information for states' internal purposes. It should not be used to compare one state with another.

To protect people's privacy, we do not show the data when 20 or fewer people from a state or residence type responded to a question (though their responses are included in the NCI-IDD average).

Important note on Impact of COVID-19

The 2020-21 NCI-IDD IPS data collection cycle began July 1, 2020 and ended June 30, 2021. As COVID-19

continued to spread across the United States, states were impacted in various ways throughout the year. These data cannot fully assess what impact the pandemic had on data collected at different times throughout the year. While these data will serve as an important baseline, and meaningful way for states to understand the overall impact of services on people being served, these data should be read with caution.

To help better understand the overall impact of COVID-19 on people receiving services, NCI-IDD added a state-optional COVID-19 Supplement to all NCI-IDD Surveys. The COVID-19 Supplement is intended to support the understanding of the experience of people with disabilities and their families at the beginning of the COVID-19 pandemic, and their interactions with services during that time. The NCI IPS COVID-19 Supplement can be found [here](#).

Figure A. Percentage of people surveyed in each residence type by state

State	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home	N
AL	0%	76%	6%	18%	0%	391
AR	1%	14%	39%	45%	2%	397
CA ¹	10%	25%	13%	51%	2%	8,630
CO	0%	11%	9%	52%	27%	614
FL	0%	42%	16%	42%	0%	399
GA	0%	31%	15%	40%	14%	405
HI	0%	13%	6%	52%	29%	317
IL	0%	50%	10%	39%	1%	339
IN	0%	27%	28%	43%	2%	405
KS	1%	43%	25%	28%	3%	380
KY	1%	37%	7%	28%	27%	393
MI	0%	48%	15%	35%	3%	651
MN	1%	55%	7%	33%	4%	386
MO	1%	20%	45%	30%	4%	399
NE	8%	29%	21%	26%	16%	417
NV	0%	51%	22%	22%	5%	408
NH	2%	15%	10%	39%	33%	396
NJ	0%	48%	13%	34%	5%	406
NC	14%	14%	8%	50%	14%	383
OH	6%	12%	36%	44%	1%	329
OR	0%	21%	9%	50%	20%	419
PA	8%	40%	12%	39%	1%	605
SD	0%	53%	36%	10%	1%	398
TN	3%	39%	31%	24%	4%	317
VA	0%	44%	8%	30%	18%	718
WI	1%	22%	22%	54%	1%	739
Total	5%	30%	16%	43%	6%	19,641

¹ All Californians with an intellectual or developmental disability as defined by California law have both a civil right and an individual entitlement to receive services from the California Department of Developmental Services. These statutory requirements make California's service system unique and could impact comparisons between its survey results and the results of other states.




Table 1. Takes at least one medication for mood, anxiety, psychotic disorder and/or behavioral challenge (information may have been obtained through state records) ²**Significantly Above Weighted NCI Average**

State	Average Within State	N	ICF/IID. Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
TN	71%	303	n/a	75%	78%	51%	n/a
KY	67%	371	n/a	74%	68%	41%	84%
KS	65%	376	n/a	71%	74%	49%	n/a
MO	64%	396	n/a	68%	73%	47%	n/a
VA	63%	685	n/a	70%	66%	51%	65%
SD	62%	381	n/a	72%	54%	34%	n/a
WI*	62%	400	n/a	78%	61%	43%	n/a
NH	62%	389	n/a	71%	49%	45%	80%
MI	60%	628	n/a	72%	60%	45%	n/a
AL	60%	381	n/a	66%	48%	34%	n/a
IL	60%	326	n/a	68%	53%	50%	n/a
PA	59%	619	50%	74%	54%	46%	n/a
IN	58%	397	n/a	82%	60%	42%	n/a
NJ	57%	392	n/a	70%	51%	46%	36%
NC	57%	373	56%	73%	63%	48%	74%

Within Weighted NCI Average

State	Average Within State	N	ICF/IID. Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
GA	55%	402	n/a	74%	56%	37%	64%
FL	53%	391	n/a	72%	47%	37%	n/a
CO	53%	559	n/a	66%	46%	39%	73%
NE	51%	394	81%	50%	44%	44%	58%
Weighted NCI-IDD Average	50%	18,408	50%	69%	53%	33%	65%
NV	50%	397	n/a	60%	51%	23%	n/a
OR	49%	410	n/a	73%	49%	34%	62%
OH*	49%	164	n/a	66%	62%	24%	n/a
AR	48%	390	n/a	57%	61%	34%	n/a

Significantly Below Weighted NCI Average

State	Average Within State	N	ICF/IID. Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
HI	35%	302	n/a	38%	n/a	25%	52%
CA	34%	8,582	44%	60%	32%	18%	44%

² Data were not available in records for MN

State data with at least 25% "don't knows" and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C




Table 2. Takes medication for mood, anxiety, and/or psychotic disorders (information may have been obtained through state records) ³**Significantly Above Weighted NCI Average**

State	Average Within State	N	ICF/IID. Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
TN	69%	301	n/a	74%	76%	48%	n/a
KY	64%	367	n/a	72%	67%	36%	83%
KS	62%	378	n/a	69%	72%	43%	n/a
MO	60%	397	n/a	61%	70%	42%	n/a
SD	60%	383	n/a	70%	50%	31%	n/a
NH	58%	393	n/a	69%	49%	40%	75%
WI*	58%	402	n/a	72%	58%	39%	n/a
VA	58%	685	n/a	64%	66%	42%	63%
PA	56%	617	45%	70%	51%	42%	n/a
IL	56%	328	n/a	64%	53%	45%	n/a
MI	56%	633	n/a	67%	56%	40%	n/a
AL	55%	382	n/a	62%	45%	30%	n/a
IN	54%	399	n/a	77%	59%	36%	n/a

Within Weighted NCI Average

State	Average Within State	N	ICF/IID. Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
NJ	52%	390	n/a	62%	51%	44%	23%
NC	52%	376	47%	71%	59%	43%	67%
GA	51%	403	n/a	70%	52%	33%	61%
CO	48%	562	n/a	62%	41%	33%	71%
OH*	48%	163	n/a	66%	62%	21%	n/a
NV	47%	400	n/a	57%	50%	20%	n/a
FL	47%	394	n/a	63%	43%	32%	n/a
Weighted NCI-IDD Average	47%	18,445	48%	65%	51%	29%	61%
OR	46%	412	n/a	72%	49%	29%	58%
AR	44%	393	n/a	54%	57%	30%	n/a

Significantly Below Weighted NCI Average

State	Average Within State	N	ICF/IID. Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
NE	38%	396	78%	38%	37%	28%	39%
CA	32%	8,589	43%	59%	30%	15%	41%
HI	26%	302	n/a	30%	n/a	17%	38%

³ Data were not available in records for MN

State data with at least 25% "don't knows" and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C

Table 2B. Takes medication for mood, anxiety, and/or psychotic disorders (*includes missing and “don’t know” responses*; information may have been obtained through state records)

State	Takes medication	Does not take medication	Missing	Don't know	N
AL	54%	44%	1%	1%	391
AR	43%	54%	1%	2%	406
CA	32%	68%	1%	0%	8,663
CO	41%	45%	5%	9%	655
FL	47%	52%	1%	0%	400
GA	51%	49%	0%	0%	405
HI	24%	70%	3%	2%	319
IL	53%	42%	3%	2%	346
IN	53%	45%	1%	1%	407
KS	61%	36%	2%	1%	390
KY	59%	33%	2%	6%	399
MI	53%	42%	1%	4%	664
MN	0%	0%	100%	0%	406
MO	59%	40%	0%	1%	402
NE	36%	58%	2%	3%	419
NV	46%	52%	1%	0%	408
NH	57%	41%	1%	0%	399
NJ	50%	46%	1%	3%	407
NC	45%	42%	12%	1%	432
OH	23%	26%	44%	7%	333
OR	45%	53%	2%	0%	420
PA	52%	41%	6%	2%	668
SD	56%	38%	3%	2%	405
TN	65%	29%	1%	5%	322
VA	55%	40%	0%	5%	719
WI	29%	21%	0%	50%	806
Extended weighted NCI-IDD Average	40%	46%	10%	5%	19,991

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C




Table 3. Number of medications taken for at least one of the following: mood disorders, anxiety, or psychotic disorders if taking at least one medication for this purpose (information may have been obtained through state records) ⁴

State	1-2 medications	3-4 medications	5-10 medications	11 or more medications	N
AL	81%	17%	2%	0%	210
AR	67%	28%	5%	0%	169
CO	75%	19%	6%	0%	262
FL	76%	15%	9%	1%	185
GA	67%	26%	7%	0%	206
HI	75%	19%	5%	0%	77
IL	73%	23%	4%	0%	181
IN	72%	20%	7%	1%	209
KS	63%	26%	12%	0%	232
KY	69%	23%	8%	0%	213
MI	59%	32%	9%	0%	346
MO	67%	22%	10%	0%	229
NE	80%	14%	5%	1%	141
NV	72%	26%	2%	0%	184
NH	60%	30%	9%	0%	223
NJ	65%	31%	4%	0%	201
NC	64%	27%	9%	1%	192
OH	73%	24%	3%	0%	67
OR	79%	19%	2%	0%	175
PA	61%	31%	8%	0%	333
SD	59%	32%	9%	0%	220
TN	70%	28%	2%	0%	164
VA	57%	33%	9%	1%	385
WI	59%	30%	11%	0%	228
Weighted NCI-IDD Average	67%	25%	7%	0%	5,032

⁴ Data were not available in records for MN and CA

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C

Table 3B. Number of medications taken for at least one of the following: mood disorders, anxiety, or psychotic disorders if taking at least one medication for this purpose (*includes missing and “don’t know” responses*; information may have been obtained through state records)

State	1-2 medications	3-4 medications	5-10 medications	11 or more medications	Missing	Don't know	N
AL	81%	17%	2%	0%	0%	0%	211
AR	65%	27%	5%	0%	3%	0%	174
CA	3%	2%	0%	0%	95%	0%	2,618
CO	73%	18%	6%	0%	3%	0%	269
FL	76%	15%	9%	1%	1%	0%	186
GA	67%	26%	7%	0%	0%	0%	206
HI	74%	19%	5%	0%	1%	0%	78
IL	72%	22%	4%	0%	1%	0%	183
IN	70%	19%	6%	1%	4%	0%	217
KS	61%	25%	11%	0%	2%	0%	236
KY	62%	20%	8%	0%	10%	0%	236
MI	58%	31%	9%	0%	2%	0%	353
MN	0%	0%	0%	0%	100%	0%	406
MO	65%	22%	10%	0%	3%	0%	237
NE	74%	13%	5%	1%	7%	0%	152
NV	70%	25%	2%	0%	3%	0%	189
NH	59%	30%	9%	0%	2%	0%	228
NJ	64%	31%	4%	0%	1%	0%	203
NC	63%	27%	9%	1%	2%	0%	195
OH	63%	21%	3%	0%	14%	0%	78
OR	73%	18%	2%	0%	7%	0%	188
PA	59%	30%	8%	0%	3%	0%	345
SD	57%	31%	8%	0%	4%	0%	228
TN	55%	22%	2%	0%	22%	0%	209
VA	56%	32%	9%	1%	3%	0%	395
WI	58%	29%	11%	0%	2%	0%	233
Extended weighted NCI-IDD Average	52%	20%	6%	0%	23%	0%	7,847

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C




Table 4. Takes medication for behavior challenges (information may have been obtained through state records) ⁵**Significantly Above Weighted NCI Average**

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
VA	39%	670	n/a	45%	22%	34%	39%
WI*	38%	394	n/a	56%	31%	24%	n/a
KS	35%	373	n/a	44%	35%	23%	n/a
TN	33%	278	n/a	30%	48%	16%	n/a
NE	32%	390	48%	30%	22%	28%	45%
NC	31%	363	43%	32%	23%	26%	40%
IL	30%	324	n/a	37%	13%	25%	n/a
AL	30%	384	n/a	37%	14%	7%	n/a
FL	29%	392	n/a	47%	12%	17%	n/a
AR	28%	390	n/a	31%	36%	22%	n/a
IN	27%	392	n/a	43%	23%	19%	n/a
KY	27%	374	n/a	34%	32%	23%	20%
NH	27%	387	n/a	31%	24%	19%	35%
NJ	27%	391	n/a	37%	12%	20%	14%
MI	25%	620	n/a	31%	18%	22%	n/a

Within Weighted NCI Average

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
MO	22%	397	n/a	25%	24%	18%	n/a
PA	22%	613	26%	30%	13%	13%	n/a
CO	21%	526	n/a	31%	14%	17%	29%
Weighted NCI-IDD Average	20%	18,313	15%	29%	17%	13%	28%
GA	19%	400	n/a	27%	20%	10%	27%
OH*	18%	165	n/a	23%	19%	9%	n/a
SD	17%	385	n/a	22%	14%	8%	n/a
OR	17%	409	n/a	22%	8%	12%	29%
NV	17%	392	n/a	21%	15%	6%	n/a
HI	15%	305	n/a	20%	n/a	8%	29%
CA	6%	8,599	5%	9%	5%	5%	7%

⁵ Data were not available in records for MN

State data with at least 25% "don't knows" and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C

Table 4B. Takes medication for behavior challenges (*includes missing and “don’t know” responses*; information may have been obtained through state records)

State	Takes medication	Does not take medication	Don't know	Missing	N
AL	30%	69%	2%	0%	391
AR	27%	69%	0%	4%	406
CA	6%	93%	1%	0%	8,663
CO	17%	63%	5%	15%	655
FL	29%	69%	2%	1%	400
GA	19%	80%	0%	1%	405
HI	15%	81%	3%	2%	319
IL	28%	65%	4%	3%	346
IN	26%	70%	1%	3%	407
KS	34%	62%	2%	3%	390
KY	25%	69%	1%	5%	399
MI	24%	70%	1%	5%	664
MN	0%	0%	100%	0%	406
MO	22%	77%	0%	1%	402
NE	29%	64%	2%	5%	419
NV	16%	80%	2%	2%	408
NH	26%	71%	2%	1%	399
NJ	26%	71%	1%	3%	407
NC	26%	58%	12%	4%	432
OH	9%	41%	43%	7%	333
OR	16%	81%	3%	0%	420
PA	20%	72%	5%	3%	668
SD	17%	79%	2%	3%	405
TN	28%	58%	2%	12%	322
VA	36%	57%	0%	7%	719
WI	19%	30%	0%	51%	806
Extended weighted NCI-IDD Average	17%	68%	9%	5%	19,991

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C




Table 5. Number of medications taken for behavior challenges if taking at least one for this purpose (information may have been obtained through state records) ⁶

State	1-2 medications	3-4 medications	5-10 medications	11 or more medications	N
AL	82%	14%	4%	0%	112
AR	75%	24%	1%	0%	102
CO	83%	14%	4%	0%	109
FL	77%	19%	4%	0%	113
GA	86%	13%	1%	0%	77
HI	74%	26%	0%	0%	46
IL	82%	16%	2%	0%	96
IN	75%	20%	5%	1%	102
KS	76%	17%	6%	0%	127
KY	76%	15%	9%	0%	89
MI	76%	17%	6%	1%	143
MO	89%	6%	6%	0%	87
NE	71%	26%	3%	0%	120
NV	86%	14%	0%	0%	65
NH	81%	16%	3%	0%	100
NJ	72%	24%	4%	0%	101
NC	74%	17%	8%	1%	108
OH	85%	15%	0%	0%	27
OR	86%	11%	3%	0%	64
PA	74%	23%	3%	0%	126
SD	83%	15%	2%	0%	66
TN	88%	10%	1%	0%	69
VA	68%	23%	8%	1%	248
WI	66%	28%	6%	0%	149
Weighted NCI-IDD Average	77%	19%	4%	0%	2,446

⁶ Data were not available in records for MN and CA

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C

Table 5B. Number of medications taken for behavior challenges if taking at least one for this purpose (*includes missing and “don’t know” responses*; information may have been obtained through state records)

State	1-2 medications	3-4 medications	5-10 medications	11 or more medications	Missing	Don't know	N
AL	79%	14%	3%	0%	3%	0%	116
AR	70%	22%	1%	0%	7%	0%	110
CA	6%	3%	0%	0%	91%	0%	500
CO	80%	13%	4%	0%	4%	0%	113
FL	76%	19%	4%	0%	1%	0%	114
GA	86%	13%	1%	0%	0%	0%	77
HI	72%	26%	0%	0%	2%	0%	47
IL	81%	15%	2%	0%	2%	0%	98
IN	72%	19%	5%	1%	4%	0%	106
KS	73%	17%	6%	0%	4%	0%	132
KY	68%	13%	8%	0%	11%	0%	100
MI	69%	16%	6%	1%	9%	0%	157
MN	0%	0%	0%	0%	100%	0%	406
MO	88%	6%	6%	0%	1%	0%	88
NE	69%	25%	3%	0%	2%	0%	123
NV	86%	14%	0%	0%	0%	0%	65
NH	79%	16%	3%	0%	3%	0%	103
NJ	70%	23%	4%	0%	3%	0%	104
NC	72%	16%	8%	1%	3%	0%	111
OH	79%	14%	0%	0%	7%	0%	29
OR	81%	10%	3%	0%	6%	0%	68
PA	70%	22%	3%	0%	5%	0%	132
SD	82%	15%	1%	0%	1%	0%	67
TN	67%	8%	1%	0%	24%	0%	91
VA	65%	22%	7%	1%	4%	0%	259
WI	65%	28%	6%	0%	1%	0%	150
Extended weighted NCI-IDD Average	67%	17%	4%	0%	12%	0%	3,060

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C




Table 6. Has a behavior plan (information may have been obtained through state records) ⁷**Significantly Above Weighted NCI Average**

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
KS	62%	383	n/a	76%	64%	38%	n/a
IN	50%	406	n/a	81%	51%	29%	n/a
NV	45%	401	n/a	63%	32%	17%	n/a
IL	45%	325	n/a	66%	28%	18%	n/a
PA	36%	631	48%	55%	16%	16%	n/a
KY	33%	396	n/a	44%	50%	17%	29%

Within Weighted NCI Average

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
CO	31%	557	n/a	45%	25%	18%	53%
NE	31%	376	91%	31%	16%	17%	38%
AR	29%	384	n/a	34%	39%	19%	n/a
SD	27%	395	n/a	41%	15%	3%	n/a
AL	26%	389	n/a	34%	9%	0%	n/a
Weighted NCI-IDD Average	26%	9,818	53%	41%	20%	12%	31%
OR	26%	416	n/a	36%	13%	20%	36%
NC	25%	378	58%	42%	18%	10%	33%
VA	23%	685	n/a	30%	14%	12%	28%
FL	21%	396	n/a	39%	13%	7%	n/a
NH	21%	396	n/a	47%	7%	5%	32%
TN	20%	311	n/a	16%	36%	5%	n/a

Significantly Below Weighted NCI Average

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
MI	20%	651	n/a	31%	13%	9%	n/a
HI*	18%	156	n/a	16%	n/a	12%	27%
GA	16%	404	n/a	29%	18%	6%	13%
WI*	15%	405	n/a	31%	4%	3%	n/a
NJ	13%	395	n/a	22%	6%	6%	n/a
MO	12%	400	n/a	14%	16%	5%	n/a
OH*	7%	182	n/a	13%	7%	0%	n/a

⁷ Data were not available in records for MN and CA

State data with at least 25% "don't knows" and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C

Table 6B. Has a behavior plan (includes missing and “don’t know” responses; information may have been obtained through state records) ⁸

State	Has behavior plan	No behavior plan	Missing	Don't know	N
AL	26%	73%	0%	1%	391
AR	28%	67%	0%	5%	406
CA	1%	4%	95%	0%	8,663
CO	26%	59%	5%	10%	655
FL	21%	78%	1%	0%	400
GA	16%	84%	0%	0%	405
HI	9%	40%	49%	2%	319
IL	42%	52%	3%	3%	346
IN	50%	50%	0%	0%	407
KS	61%	37%	2%	0%	390
KY	33%	66%	0%	1%	399
MI	20%	78%	1%	1%	664
MN	0%	0%	100%	0%	406
MO	12%	88%	0%	0%	402
NE	27%	62%	3%	7%	419
NV	44%	54%	1%	1%	408
NH	21%	78%	0%	1%	399
NJ	13%	84%	1%	2%	407
NC	22%	66%	11%	1%	432
OH	4%	51%	42%	3%	333
OR	26%	73%	1%	0%	420
PA	34%	61%	4%	1%	668
SD	27%	71%	1%	1%	405
TN	20%	77%	1%	2%	322
VA	22%	73%	0%	5%	719
WI	7%	43%	0%	50%	806
Extended weighted NCI-IDD Average	16%	46%	34%	4%	19,991

⁸ Data were not available in records for MN and NY

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C




Table 7. Has a behavior plan (among those who take medication for a behavior challenge; information may have been obtained through state records)⁹**Significantly Above Weighted NCI Average**

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
KS	89%	132	n/a	96%	82%	78%	n/a
NV	85%	65	n/a	86%	n/a	n/a	n/a
IN	82%	106	n/a	93%	88%	61%	n/a
IL	79%	94	n/a	93%	n/a	42%	n/a
NE	73%	111	n/a	76%	n/a	54%	81%
AL	72%	116	n/a	76%	n/a	n/a	n/a
AR	72%	99	n/a	n/a	72%	65%	n/a
PA	72%	130	n/a	83%	n/a	38%	n/a

Within Weighted NCI Average

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
OR	69%	68	n/a	n/a	n/a	75%	59%
KY	67%	100	n/a	77%	n/a	50%	n/a
SD	62%	66	n/a	71%	n/a	n/a	n/a
CO	60%	101	n/a	n/a	n/a	47%	79%
FL	57%	113	n/a	70%	n/a	24%	n/a
Weighted NCI-IDD Average	56%	2,494	82%	67%	46%	35%	57%
NC	53%	109	n/a	n/a	n/a	32%	62%
MI	52%	154	n/a	64%	n/a	31%	n/a
NH	50%	103	n/a	n/a	n/a	21%	62%
TN	44%	89	n/a	40%	58%	n/a	n/a

Significantly Below Weighted NCI Average

State	Average Within State	N	ICF/IID, Nursing Facility and Other Institutional Settings	Community-based Group Residential Settings	Own Home or Apartment	Parent's or Relative's Home	Foster Care or Host Home
VA	43%	254	n/a	54%	n/a	28%	39%
GA	41%	76	n/a	53%	n/a	n/a	n/a
NJ	35%	102	n/a	41%	n/a	25%	n/a
HI	30%	43	n/a	n/a	n/a	n/a	42%
WI	30%	149	n/a	45%	11%	3%	n/a
MO	29%	86	n/a	n/a	37%	14%	n/a
OH	25%	28	n/a	n/a	n/a	n/a	n/a

⁹ Data were not available in records for MN and CA

State data with at least 25% "don't knows" and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C

Table 6B. Has a behavior plan among those who take medication for a behavior challenge (includes missing and “don’t know” responses; information may have been obtained through state records) ¹⁰

State	Has behavior plan	No behavior plan	Missing	Don't know	N
AL	72%	28%	0%	0%	116
AR	65%	25%	0%	10%	110
CA	8%	2%	90%	0%	500
CO	54%	35%	0%	11%	113
FL	56%	43%	1%	0%	114
GA	40%	58%	1%	0%	77
HI	28%	64%	4%	4%	47
IL	76%	20%	1%	3%	98
IN	82%	18%	0%	0%	106
KS	89%	11%	0%	0%	132
KY	67%	33%	0%	0%	100
MI	51%	47%	1%	1%	157
MN	0%	0%	100%	0%	406
MO	28%	69%	0%	2%	88
NE	66%	24%	1%	9%	123
NV	85%	15%	0%	0%	65
NH	50%	50%	0%	0%	103
NJ	35%	63%	0%	2%	104
NC	52%	46%	0%	2%	111
OH	24%	72%	3%	0%	29
OR	69%	31%	0%	0%	68
PA	70%	28%	2%	0%	132
SD	61%	37%	0%	1%	67
TN	43%	55%	0%	2%	91
VA	42%	56%	0%	2%	259
WI	30%	69%	0%	1%	150
Extended weighted NCI-IDD Average	51%	40%	9%	1%	3,060

¹⁰ Data were not available in records for MN

State data with at least 25% “don’t knows” and missing data are denoted with an asterisk (*)

Data for WI were unable to be collected for those respondents who self-direct their services

See all un-collapsed state data showing responses for all possible response options in Appendix C