

Exploring Health Disparities Among People with Intellectual and Developmental Disabilities

What Are the Issues and Do Race and Ethnicity Play a Role?



Valerie J. Bradley, Human Services Research Institute



NASDDDS Annual Meeting November 13, 2014

Agenda

Today...
1.
2.
3.

- Research questions
- What are health disparities?
- Health/healthcare disparities and people with ID/DD
- What do NCI data show?

Research Questions


- Do people with intellectual and developmental disabilities experience health disparities overall?
- Do NCI data demonstrate differences in utilization and access to preventive healthcare by race/ethnicity?
- Do NCI data demonstrate differences by race/ethnicity in other outcomes, such as employment, choice, community inclusion, and rights/respect?

National Core Indicators (NCI)?



- NASDDDS HSRI Collaboration
- Multi-state collaboration of state DD agencies
- Measures performance of public systems for people with intellectual and developmental disabilities
- Assesses performance in several areas, including: employment, community inclusion, choice, rights, and health and safety
- Launched in 1997 in 13 participating states now in 42 states (including DC) and 22 sub-state areas
- Now expanded to elderly and people with disabilities through the NCI-AD

NCI State Participation 2014-15



■ State contract awarded in 2014-15 through AIDD funding CA*- Includes 21 Regional Centers

OH*- Also includes the Mid-East Ohio Regional Council


How Does NCI Collect Data?

- Adult Consumer Survey
 - ✓ In-person conversation with a sample of adults receiving services to gather information about their experiences
 - ✓ Keyed to important person-centered outcomes that measure system-level indicators related to: employment, choice, relationships, case management, inclusion, health, etc.
- Adult Family, Child Family, and
 Family/Guardian Surveys: mail surveys –
 separate sample from Adult Consumer Survey
- Other NCI state level data: Mortality, Staff Stability





What Are Health Disparities?



Health Disparity Populations

The National Institute on Minority Health and Health Disparities (NIMHD) defines the population as those with:

A significant disparity in the overall rate of disease incidence, prevalence, morbidity, mortality, or survival rates in the population as compared to the health status of the general population.

Health Disparity Populations


Current NIMHD health disparity target groups include:

- Blacks/African Americans
- Hispanics/Latinos
- American Indians/Alaska Natives
- Asian Americans
- Native Hawaiians and other Pacific Islanders
- Socioeconomically disadvantaged populations and rural populations
- Disability advocates pressing for inclusion in the definition



Health Determinants

- Why are some people healthy/unhealthy?
- Differences in health/
 healthcare utilization exist
 because of unequal
 distributions of social,
 environmental, economic
 conditions within society
- NCI captures demographics, employment, communication, choice., etc.



Health Determinants Model

 $http://www.health-inequalities.eu/HEALTHEQUITY/EN/about_hi/health_inequalities/$




Do Individuals with ID/DD Experience Health/Healthcare Disparities?

ID/DD Population as Potential Health Disparity Population?

 American Academy of Developmental Medicine and Dentistry (AADMD), AMA, and ADA:

Advocating to have ID/DD population designated as "medically underserved" by Health Resources and Services Administration—would lead to increase in resources to address disparities

- People with ID/DD shown to be "socioeconomically disadvantaged*"
- People with ID/DD experience:
 - Higher rates of certain diseases/conditions
 - More deaths and morbidity from those diseases when compared with the general population



*Rehabilitation Research and Training Center on Disability Statistics and Demographics (StatsRRTC), 2004 Disability Status Reports 2004; Ithaca (NY): Cornell University



What Are the Top 5 Causes of Death in the General Population?

Rank		Cause		
1		Heart Disease 24.5%		
2		Cancer 23.3%		
3		Emphysema, Asthma, 5.68%	Bronchitis	
4		Stroke 5.3%		
5		Unintentional Injury	5.02%	

http://www.cdc.gov/nchs/fastats/leading-causes-of-death.htm



What Do People with ID/DD Die of?

Comparison of the Top 5 Leading Causes of Death as Reported by Four State ID/DD Agencies

Rank	MA DDS	MA DDS	CT DDS45	OH DDD47	LA OCDD
	CY2010	CY2011	FY2011	2010	FY2011
	(Adults)	(Adults)	(all ages)	(Adults)	(all ages)
Method	Underlying		Primary	Unknown	Unknown
1	Heart Disease	Heart Disease	Heart Disease	Heart Disease	Heart Disease
	18.0%	17.5%	27.4%	18.9%	18.4%
2	Cancer	Cancer	Cancer	Cancer	Septicemia
	13.8%	12.7%	13.5%	11.3%	17.3%
3	Alzheimer's	Alzheimer's	Aspiration	Influenza &	Congenital
	Disease	Disease	Pneumonia	Pneumonia	Condition
	12.8%	10.9%	12%	9.0%	16.3%
4	Aspiration Pneumonia 12.3%	Aspiration Pneumonia 8.0%	Respiratory Disease4 11.1%	Congenital Diseases 8.3%	Pneumonia 10.2%
5	Septicemia 8.6%	Septicemia 7.7%	Pneumonia 8.2%	Aspiration Pneumonia 7.1%	Malignant Neoplasm (Cancer) 7.1%

At What Ages Do People with ID/DD Die?

Average Age at Death by Gender, 2011

Gender	Average Age at Death MA DDS	Average Age of Death US General Population	
Female	62.5	81.1	
Male	59.9	76.3	

What Does the CDC Say About the Health Status of People with ID/DD?

Adults with intellectual disabilities experience poorer health outcomes than people without ID. These disparities mean that people with ID are more likely to:

- Live with complex health conditions.
- Have limited access to quality healthcare and health promotion programs.
- Miss cancer screenings.
- Have poorly managed chronic conditions, such as epilepsy.
- Be obese.
- Have undetected poor vision.
- Have mental health problems and use psychotropic medications.

http://www.cdc.gov/ncbddd/developmentaldisabilities/index.html



ID/DD Healthcare Disparities in Preventive Care Use

Women with ID/DD are less likely than women without ID/DD to:

- Have had cervical and breast cancer screenings
- Have ever visited a gynecologist

Individuals with ID/DD are less likely than individuals without ID/DD to:

- Visit dentist regularly
- Get eye and hearing tests
- Receive timely vaccines





Do Ethnic and Racial
Disparities Exist Regarding
Utilization of Preventive
Care?

What Is Preventive Care?

- Prevents people from getting sick by detecting diseases/conditions before they become serious
- Beneficial for both financial purposes and to achieve a high quality of life
- Enhanced attention to preventive care:
 - Patient Protection and Affordable Care Act (ACA)
 - Requires that many health insurance companies cover the full cost to the consumer of many types of regular preventive care services (Healthcare.gov).

Racial/Ethnic Healthcare Disparities in Use of Preventive Care

General Public:

African American and Hispanic individuals

- Visit their personal physician and have dentist visits less frequently than whites
- Are less likely than whites to receive services such as
 - Flu and/or pneumonia vaccines
 - Colorectal cancer screenings
 - Pap tests
 - Mammograms





What Do NCI Adult Consumer Data Tell Us About Racial and Ethnic Disparities in Utilization of Preventive Care?

Data Source: Adult Consumer Survey

- Standardized, face-to-face interview with a sample of individuals receiving services
 - Background Information includes health information
 - Section I (no proxies allowed)
 - Section II (proxies allowed)
- No pre-screening procedures
- Conducted with adults only (18 and over) receiving at least one service in addition to case management
- Section I and Section II together take 50 minutes (on average)





General Findings




- Individuals living in structured settings
 (i.e., specialized institutions, group
 homes) are more likely to have access to
 preventive screenings and
 vaccinations—regardless of their race
 or ethnicity.
- Individuals living with their families and those living independently are less likely to receive preventive care.

Residence Type


- To determine whether race or ethnicity played a role in access to preventive care, we only included individuals living in:
 - Independent home/apartment
 - Parent/relative's home
 - Foster care/host home
- N=7,632 from 25 states
- Race/ethnicity collapsed into 3 groups African American (non Hispanic), Hispanic (including black Hispanic), White (non-Hispanic)

Demographic Differences

Final Sample Residence Type (p<=.001)



Race/Ethnicity of Individuals Living at Home, Independently or in a Foster Home/Host Home



Preventive Care Utilization Disparities

White respondents, African American respondents, and Hispanic respondents have significantly different rates of:



- Having had a physical exam in the past year (p<.001)
- Having gone to the dentist in the past year (p<.001)
- Having had a routine vision screening/eye exam in the past year (p<.001)
- Having had a flu vaccine in the past year (p<.001)
- Having ever had a pneumonia vaccine (p<.001)


Important to Control for Other Factors

- Individuals of different races and ethnicities also differ in other demographic characteristics such as:
 - State of residence
 - Age
 - Individual's primary language
 - Individual's primary means of expression

- Level of intellectual disability
- Mobility
- Other diagnoses (in addition to ID/DD)
- Residence type
- These differences may affect their rates of preventive care use.
- Some examples of demographic differences are seen in the next few slides...




Level of ID by Race/Ethnicity



- White respondents more likely to be diagnosed with mild ID
- African American and Hispanic respondents more likely than White respondents to be diagnosed with moderate, severe, or profound ID


Mobility Level by Race and Ethnicity





Other Diagnoses by Race/Ethnicity

Mental Illness/Psychiatric Diagnosis



- White respondents are more likely to be diagnosed with mood disorder and/or anxiety disorder
- African American and Hispanic respondents more likely to be diagnosed with psychotic disorder

After controlling for demographic factors...

For individuals not living in institutional or community-based settings, the following preventive care exams showed differences by race/ethnicity that were still statistically significant:



African American respondents and Hispanic respondents were less likely than White, Non-Hispanic respondents to have had a physical exam in the past year.




African Americans were less likely than Whites to have had a dentist visit in the past year.



African Americans were less likely than Whites to have had a flu vaccine in the past year.

Example:

Though Whites Are More Likely to Have Ever Gotten
Pneumonia Vaccine (p < .001) – Bigger Predictor was Risk
Factors Including Age (older), Down Syndrome, Poor Health,
and Reduced Mobility





What Do NCI Adult Family
Survey Data Tell Us About
Racial and Ethnic Disparities
in Access to Preventive Care?

Adult Family Survey 2012-13

- Mail-in surveys
- Adult Family Survey: respondent is a family member of an individual with ID/DD over age 18 who lives in the family home
- 2012-13: N=5,010
- 13 states






Administration of Adult Family Survey

- Mail-in
 - Selection bias!
- Sent to families with an adult family member living at home
- The Adult Family Survey (AFS) asks about *access* to health services as opposed to *utilization*. For example:
 - "Do you have access to health services for your family member."
- Also asks about dental services, medications, and mental health services.



Demographic Breakdown

Adult Family Survey 2012-13 (N=4,760*)



*Only those cases for which a racial/ethnic identity was provided were included in this analysis.




Though Data Showed Some Differences in Access (e.g., dental, doctors visits) Needed to Control For:

- State of residence
- Family member's age
- Whether there is more than one person with a disability in the household
- Family member's primary language
- Family member's primary means of expression
- Family member's other diagnoses (in addition to ID/DD)

- Family member's highest level of education
- Frequency at which family member requires medical care
- Whether family member needs behavior support
- Level of support family member needs for activities of daily living
- Respondent's highest education level
- Household income

For example...


More than One Person With ID/DD Lives in Household (p<=.001)



Hispanic respondents significantly more likely to need moderate or complete help with daily activities (such as bathing, dressing, eating) than White, Non-Hispanic and African American, Non-Hispanic respondents.

For example.....

Household Income in Past Year (p<=.001)



Findings

Interestingly...

 Household income is a significant predictor of access to dental care.

When we control for these demographic differences

 Race/Ethnicity is no longer a significant predictor of access to any of the preventive care specified in the AFS.



Do NCI Data Demonstrate Differences by Race/Ethnicity in Other Valued Outcomes?

Using full NCI ACS dataset: N=13,157

Other Outcome Disparities



There are significant differences in rates of paid community employment among the racial and ethnic categories.

In addition, among those who don't have a paid community job, African Americans more likely to want a paid community job.

White respondents were significantly more likely to have had at least some input in:

Choosing home, schedule, what to do in free time, and/or choosing day activity

Other Outcome Disparities (cont.)

Hispanic respondents are more likely to report that:



- Their email/mail is read without their permission
- There are rules against being alone with friends/visitors
- There are rules/restrictions on use of phone/internet

Summary

- While people with ID/DD do experience health disparities and are disproportionately affected by disease, and untimely death, they are not a federally designated group.
- People with ID/DD are more likely to die of preventable causes and to die at an earlier age.



Summary (cont.)

While a variety of factors may explain disparate access to preventive health services,



African American respondents and Hispanic respondents were less likely than White respondents to have had a physical exam in the past year



African Americans were less likely than Whites to have had a dentist visit in the past year



African Americans were less likely than Whites to have had a flu vaccine in the past year

Summary (cont.)



 Family income is a bigger predictor than race and ethnicity of access to preventive health services.



 NCI data show that African Americans and Hispanic respondents to the Adult Consumer Survey are less likely to achieve outcomes such as employment and choice and to have their rights respected.


What To Do Now?

- Advocate that HRSA include individuals with ID/DD as a medically underserved population.
 - Allocate more resources to research into disparities
- As more people with ID/DD live in their own homes, with their families and in foster/host homes, public managers and MCOs will need to find other means to ensure that individuals with ID/DD utilize preventive services especially in light of their health challenges
 - Track data
 - Targeted outreach campaigns to different populations





What To Do Now? (cont.)



- Conduct continued research into racial and ethnic disparities in preventive and general healthcare
 - Help tease out root causes/social determinants of disparities and what can be done to mitigate them
- Support more systematic examination of mortality data for individuals with ID/DD
- Track the implementation of the ACA including broader coverage, better training, more accessible facilities
- Explore reasons for differential results regarding employment, choice, rights

Selection of References:

- Agency for Healthcare Research and Quality. Addressing Racial and Ethnic Disparities in Health Care. April 2013. Available at http://www.ahrq.gov/research/findings/factsheets/minority/disparit/index.html . Accessed September 24, 2013.
- Ashton C, Haidet P, Paterniti D, Collins T, et al. Racial and ethnic disparities in the use of health services. Journal of General Internal Medicine 2003; 18(2):146-152
- Bershadsky J, Kane R. Place of residence affects routine dental care in the intellectually and developmentally disabled adult population on Medicaid. Health Services Res 2010: 45(5) pt 1: 1376-89
- Bershadsky J, Taub S, Bradley V, Engler J, Moseley C, Lakin KC, et al. Place of residence and preventive health care for developmental disabilities services recipients in twenty states. Public Health Reports 2012; 127, 475-485.
- Bonito A, Eicheldinger C, Lenfestey N. Health disparities: Measuring health care use and access for racial/ethnic populations. Final Report, Part 2. 2005. Available at http://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/Reports/downloads/bonito_part2.pdf. Accessed September 24, 2013
- Escarce J. Racial and ethnic disparities in access to and quality of health care. The Synthesis Project; Robert Wood Johnson Foundation 2007; Available at www.policysynthesis.org. Retrieved September 24, 2013
- Havercamp S, Scandlin D, Roth M. Health disparities among adults with developmental disabilities, adults with other disabilities, and adults not reporting disability in North Carolina. Public Health Reports 2004; 119:418-426
- Institute of Medicine. "Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care." Washington DC: National Academy Press; 2002.
- Parish, S., Swaine, J., Son, E., Luken, K. (2013) *Receipt of mammography among women with intellectual disabilities: Medical record data indicate substantial disparities for African American Women.* Disability and Health Journal. 6:36-42

For more, please contact dhiersteiner@hsri.org



Contacts

HSRI Val Bradley: vbradley@hsri.org

NASDDDS Mary Lee Fay: MLFay@nasddds.org

NCI website <u>www.nationalcoreindicators.org</u>



